

MODUIC 3 Self Monitoring for Relapse Prevention

centents

	<u>Page</u>
Self Monitoring	2
Mood Monitoring Worksheet	3
Symptom Monitoring Worksheet	4
Being Prepared	5
Early Warning Signs Worksheet	6
A Plan of Action	7
Charting Your Life	9
Module Summary	10

self menitering

How do we know that a storm is about to break? Perhaps we hear thunder in the distance, the skies are dark and grey with rain clouds, and it becomes very windy. If we do not want to be caught with the worst effects of the storm, we would probably go back to our homes, close all the windows, bring our washing in from the line, and secure any loose items outside the house. Similarly, if bipolar patients become more aware of their early warning signs that signal the onset of a mood episode, they can take steps to prevent a full-blown episode of depression and mania.

Early detection of an impending "storm" – in the case of bipolar disorder, an episode of mania or depression – can lead to early intervention and prevention of a mood episode. In order to be able to detect an oncoming episode, bipolar patients must learn to recognise their own early warning signs and symptoms of their illness. Each person has their own unique set of signs and symptoms although many will be common to other patients.

It is not enough to be able to recognise and detect early warning signs and symptoms of a mood episode, you need also to monitor them regularly. It will not be much good if you are aware of your early warning signs and symptoms, but you continue to go through your daily life without paying much attention to the onset of the symptoms. Similarly, a person who knows that ominous grey clouds, thunder, lightning, and rushing wind signal an impending storm can still get caught in the storm if they were sitting on a park bench too engrossed in a book to look around them. Therefore, regular self monitoring is important for the purpose of early intervention to prevent relapse.

Mood Monitoring



The first step to self monitoring is monitoring your mood for the day. Ask yourself, "How did I feel today? Was my mood within the normal range, or was I feeling slightly low or high? How low? How high?" Rate your mood, between -5 (depressed) and +5 (manic). Try to rate your mood at the same time everyday.

Use the worksheet on the next page to monitor your moods. Try it for a week and see if you can identify any pattern in your mood fluctuations. You might also want to take note of the circumstances in which you experienced particularly high or low moods. Bring your completed mood monitoring worksheets to your doctor or mental health professional and discuss your observations with them.

Symptom Monitoring

Another way of monitoring yourself is to identify and be aware of any signs and symptoms associated with a depressive, manic, or mixed episode that you might be experiencing. If you experience a number of these symptoms over a few days, in such a way that they interfere with most of your day-to-day activities, you might want to consider taking some action. We will talk about planning early interventions a little later on. Use the worksheet on page 4 to record any symptoms that might have interfered with





Over the next week, monitor the fluctuations in your daily mood by filling in the mood graph below.

	Day I	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Date:							
Manic							
+5	•	•	•	•	•	•	•
+4	•	•	•	•	•	•	•
+3 Time to Intervene	•	•	•	•	•	•	•
+2 Monitor Closely	•	•	•	•	•	•	•
+	•	•	•	•	•	•	•
0 Normal	•	•	•	•	•	•	•
-1	•	•	•	•	•	•	•
-2 Monitor Closely	•	•	•	•	•	•	•
-3 Time to Intervene	•	•	•	•	•	•	•
-4	•	•	•	•	•	•	•
-5	•	•	•	•	•	•	•
Depressed							
Circumstances in which	l experie	nced par	ticularly	high or l	ow mood	ls:	

symptom monitoring

Each day, use the record sheet below and tick (\checkmark) to indicate if you have experienced any of the symptoms listed in the table below, in a way that they have **interfered with most of your day-to-day activities**. You might also want to record any observations you may have about the circumstances in which you experienced these symptoms.



	Day I	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Dat	:e:						
Depressed mood							
Loss of interest or pleasure							
Increase or decrease in appetite							
Unable to sleep or sleep too much							
Physically agitated or slowed down							
Fatigue or loss of energy							
Feeling worthless or guilty							
Unable to concentrate or make decisions							
Thoughts of death or suicide							
Elevated or irritable mood							
Increased self-esteem or self-confidence							
Decreased need for sleep							
More talkative than usual							
Racing thoughts							
Easily distracted							
Increase in goal-directed activity							
Overly eager to engage in pleasurable activities							

Circumstances in which I experienced these symptoms:



Being prepared

Identifying and Monitoring Your Early Warning Signs

Early warning signs of a relapse or an episode recurrence are symptoms that typically signal the onset of a mood episode. Some patients may think that they are not able to predict an episode but researchers have found that many bipolar patients are able to recall early warning symptoms that come before a full-blown episode. Bipolar patients have commonly reported increased activity, decreased need for sleep, and elevated mood as early warning signs of mania, and depressed mood, loss of energy, loss of interest in people or activities, impaired concentration, and thoughts of death as early warning signs for depression. It was noted that although there was some differences in the specific early warning symptoms experienced by patients, these symptoms appeared to be the same within each patient in subsequent episodes. Thus, although some early warning symptoms may be unique to individual patients, it appears that they are quite accurate in predicting the onset of a mood episode for each patient.

Remember the Symptom Record in Module 1? If you completed that worksheet, you would have identified and recorded your typical symptoms of mania and depression, and distinguished those from what you are like when you are normal. To identify your early warning signs of mania and depression, a finer analysis needs to be done. Ask yourself, "What am I like when my mood is mildly elevated and moderately elevated? What am I like when I am mildly depressed and moderately depressed? Use the Early Warning Signs Worksheet on the next page to record all these symptoms. When you have written them down, reflect on the three or four most prominent early warning symptoms of mania and depression. Will you be able to recognise these symptoms when you next experience them? It might be a good idea to discuss this worksheet with your doctor or appropriate health professional.

Planning Early Interventions



The next step is to develop an action plan that details what you will do when you recognise the early warning signs and symptoms of a major depressive or manic episode. The worksheet on page 7 can be used to record your early intervention plans. It is important to be prepared so that when the time comes, you will know what to do. Plan what you will do, what you will say, what you will ask your friends and family to do for

you, etc. For example, your action plan may include a visit to your doctor when you recognise your energy level has increased and are feeling restless, or you may ask a friend to keep your credit card when you have the urge to shop for shoes, or request that a relative drop by to visit you when you stop calling them, etc. You may also want to ask your friends and family to say specific things to you to highlight the possibility that you might be becoming unwell. It's probably good to let them know the best way to say it so it doesn't offend you. Detail your early intervention plans carefully and keep them on your desk or on the refrigerator – somewhere that is easily accessible or visible – so that you can refer to them when the need arises.





warning signs worksheet

What am I like when I am mildly depressed?	What am I like when I am moderately depressed?
What am I like when my mood is mildly elevated?	What am I like when my mood is moderately elevated?




My most significant early warning signs are:

Early Warning Signs of Mania	Early Warning Signs of Depression
y plan of action for relapse prevention:	
What I would do to prevent a	What I would do to prevent a
full-blown Maric episode	full-blown depressive episode
What I would ask my friends	What I would ask my friends
or family to do for me	or family to do for me
What I would ask my friends or family	What I would ask my friends or family
to say to me	to say to me
•	
entre for	<u> </u>

charting your life

Research has shown that for many people, their first episode was triggered by a major life stressor. Subsequent episodes have also been linked with a stressful life event. Because of this, we encourage you to think about and chart the course of illness in your life by doing a Life Chart. A life chart is a diagram that illustrates the number, sequence, and duration of manic and depressive episodes, beginning from the very first episode. The purpose of drawing a life chart is for you to track and identify patterns of recurrences, early warning signs, and to see if there has been anything that might have triggered the onset of an episode. As such, it is important that you include in your life chart the occurrence of significant life events and the influence of treatment, such as medication and psychotherapy.

Below is an example of a life chart that tracks the course of illness in the life of a person, whom we will call Jane Smith. Jane, a 37-year-old woman with bipolar disorder, experienced her first episode of depression at age 20. This episode lasted almost a year and she got better without treatment. She got married and her first child, a daughter, was born soon after. This was when she had her first manic episode, but she was misdiagnosed as having a hormonal fluctuation. After her manic episode, she experienced a period of severe depression and she was treated with antidepressants by her doctor. Two years later, Jane experienced another bout of depression, which she associated with her being made redundant at work. She was treated again with antidepressants, but this time it resulted in a manic episode. She was hospitalised where she was diagnosed with bipolar disorder, and prescribed lithium. Jane found the lithium helpful in controlling her symptoms and continued taking it regularly. However, when she felt better, she realised that she had gained 10 kilos in weight and decided to stop medication. In a few weeks, she began feeling better and more energetic. She started a few craft projects but she found herself doing more and more. Soon, her symptoms were recognised as a manic episode and she was hospitalised. Several years later, she again discontinued her medication and this time she experienced dysphoria and hypomania.







To draw a life chart, draw a line in the middle of a page. This line represents a "normal" state, that is, normal compared to the times when you have felt depressed or hypomanic/manic. Next, mark the extreme right point of the line as "Today" and draw in your current mood. On the extreme left of the line, draw in your first recognisable episode of depression or hypomania/mania. Points below the line represents depressed mood and points above the line represents elevated mood. The distance from the reference line

indicates the severity of the symptoms. The length of each episode is represented by the width of each episode drawn in.

You may find it easier to begin with the most recent episode and then work backwards. Talking with family members or consulting medical records may also be beneficial. After you have completed your life chart, take a good look at all the events you have recorded. Is there any pattern of episode recurrence? Is there anything or any event that might have triggered a mood episode? Once you become more aware of the things that affect you and your moods, you can take steps to be better prepared to deal with problems or events that come your way. The ultimate goal is for you to become your own expert in dealing with and managing your illness.

Conclusion

The purpose of mood-monitoring, identifying early warning signs, and doing a life chart is to help you become more aware of yourself, your moods, and what is going on in your life so that you can manage your illness better, make informed decisions, prevent further illness episodes, and ultimately have the quality of life that you want. You may think that this makes you very focused on yourself and your illness or problems. However, by monitoring your moods and symptoms, you can be more prepared for action to prevent an episode from getting out of hand and ultimately endangering yourself. Thus, the benefits of self-monitoring far outweigh the costs.

Page 9

module sunnary

- Regular self monitoring is important for the purpose of early intervention to prevent relapse
- Mood monitoring involves rating your mood about between once a day, asking yourself, "How did I feel today? Was my mood within the normal range, or was I feeling slightly low/high? How low (0 to -5)? How high (0 to +5)?"
- Symptom monitoring is all about identifying and being aware of any signs and symptoms associated with a depressive, manic, or mixed episode that interfere with most of your day-to-day activities
- Bipolar patients commonly report increased activity, decreased need for sleep, and elevated mood as early warning signs of mania, and depressed mood, loss of energy, loss of interest in people or activities, impaired concentration, and thoughts of death as early warning signs for depression
- Identifying your early warning signs requires you to be aware of your symptoms when your mood is mildly elevated, moderately elevated, mildly depressed, and moderately depressed
- It is important for you to develop an early intervention plan that details your
 most significant early warning signs, what you would do, what you could ask
 your friends and family to do, and what you could ask them to say to you to
 warn you that you might be becoming unwell
- Stressful life events can sometimes contribute to triggering a mood episode and it would be important to be aware of how they affect you so that you can then be better prepared to deal with them
- Consistent and regular self-monitoring and early intervention are the keys to preventing relapse and episode recurrences

keep g**ein**g ...

In the next module, we will discuss a number of behavioural strategies for managing and preventing depression.

